HEALTH PROGRAM FOR DAIRY CATTLE

Group		VACCINE, DEWORMER & SUPPLEMENTS		TIME/AGE OF ADMINISTRATION	RECOMMENDED PRODUCT
		ESSENTIAL	OPTIONAL		
Replacement Calves			Heartwater	1-3 wks	Infective blood vaccine
			Salmonellosis	1 & 3 weeks	Inactivated vaccine
		Anthrax		3-6mo. Boost 4wks later	Trio-Sure OR Botu-Sure + Deca-Sure
		Botulism			
		Multi-clostridial vaccine			
		Lumpy skin disease		3-6mo	Live attenuated vaccine
			Rift Valley Fever	3-6mo	Live or Inactivated vaccine
			Pasteurellosis	3-6mo	Inactivated vaccine
			BVD/PI/IBR/BRSV	3-6mo	Inactivated/Modified live vaccine
			Redwater	3-9mo	Infective blood vaccine
			Gallsickness	3-9mo	Infective blood vaccine
		Brucellosis		4-6mo	S19 strain OR
				4-6mo & booster	RB51 strain
		Roundworms & Tapeworms		3-4 & 5-6mo	Endo + Lint
		Liver fluke &/or Conical fluke (depending on area)		3-6mo	Various-refer to catalogue
		Trace mineral supplement		3 & 6 mo	Complex A & E for cattle
Replacement heifers	Before breeding	Brucellosis		Booster 12-16 mo after initial vaccination*	*RB51 strain only
		Anthrax		Before breeding/calving (6-8wks), 12mo after calf vaccination	Trio-Sure OR Botu-Sure + Deca-Sure
		Botulism			
		Multi-clostridial vaccine			
		Lumpy skin disease		Aug to Oct	Live attenuated vaccine
		Three-day Stiff-sickness		Before breeding (6wks)	Live attenuated vaccine
			BVD/PI/IBR/BRSV	Before breeding (8wks)	Inactivated/Modified live vaccine
			Campylobacter (Vibriosis)	Before breeding (4-8wks)	Inactivated vaccine
			Trichomoniasis	Before breeding (4-8wks)	Inactivated vaccine
	Before calving		E.coli, Rota & Coronavirus	Before calving (4-8wks)	Inactivated vaccine
			Salmonellosis	Before calving (4-8wks)	Inactivated vaccine
	Strategic	Tick & Fly control			Various-refer to catalogue
		Roundworms & Tapeworms		Before 1st breeding (6wks)	Various-refer to catalogue
		Liver fluke &/or Conical fluke (depending on area)		Before 1st breeding (6wks)	Various-refer to catalogue
		Trace mineral supplement		6wks before breeding & calving	AD_3E + Complex A & E for cattle

*RB51 strain boosters: 2nd: 12-16mo & 3rd: 4w before breeding or after calving depending on breeding age.

AD_3E: Reg. No. G3796 Act 36/1947. NSO V14/19.1/1241 Act 13/2003. Botu-sure: Reg. No. G4258 Act 36/1947, NSO V18/24.4/1436 Act 13/2003. Complex A&E Cattle: Reg. No. G4120 Act 36/1947, NSO V17/19.2/1390 Act 13/2003. Deca-sure: Reg. No. G4280 Act 36/1947, NSO V19/24.4.2/1443 Act 13/2003. Endo+Lint: Reg.No. G3570 Act 36/1947. NSO V05/18.1.8/289 Act 13/2003. Trio-sure: Reg. No. G4242 Act 36/1947, NSO V18/24.4/1439 Act 13/2003.

KAH/PM051/2023/1

HEALTH PROGRAM FOR DAIRY CATTLE

Group		VACCINE, DEWORMER & SUPPLEMENTS		TIME/AGE OF ADMINISTRATION	RECOMMENDED PRODUCT
		ESSENTIAL	OPTIONAL		
Cows	Annual	Anthrax		Annually (before calving (6-8wks)	Trio-Sure OR Botu-Sure + Deca-Sure
		Botulism			
		Multi-clostridial vaccine			
		Lumpy skin disease		Aug to Oct (annually)	Live attenuated vaccine
		Three-day Stiff-sickness		Aug to Oct (annually)	Live attenuated vaccine
	Before calving / at dry off		Rift Valley Fever	Aug to Oct	Live or Inactivated vaccine
			E.coli, Rota & Coronavirus	Before calving (4-8wks)	Inactivated vaccine
			Salmonellosis	Before calving (4-8wks)	Inactivated vaccine
	After calving / before breeding		Vibriosis	Before breeding: 4-8wks	Inactivated vaccine
			Trichomoniasis	Before breeding: 4-8wks	Inactivated vaccine
			Brucellosis	Before breeding: at / close after calving	RB51 strain only
			BVD/PI/IBR/BRSV	Before breeding	Inactivated/Modified live vaccine
			Pasteurellosis	Before breeding	Inactivated vaccine
	Strategic	Tick & Fly control			Various pour-ons & concentrates - refer to catalogue
		Roundworms		As needed / According to season - see below	Levamintic Concentrate
		Liver fluke &/or Conical Fluke (depending on area)		According to season-see below	Levoxy 5.9%*Refer to package insert for withdrawals
		Trace mineral supplement		6wks before breeding & calving	AD_3E + Complex A & E for cattle
The above program is only a guideline. Speak to a / your veterinarian for more information.					
Roundworm, Liver- & Conical fluke treatment is based on the season. Autumn (Apr-May), Spring (Aug-Sep) & Mid summer (Jan-Feb) treatments are the most effective.					

AD_3E: Reg. No. G3796 Act 36/1947. NS0 V14/19.1/1241 Act 13/2003. Botu-sure: Reg. No. G4258 Act 36/1947, NS0 V18/24.4/1436 Act 13/2003. Complex A&E Cattle: Reg. No. G4120 Act 36/1947, NS0 V17/19.2/1390 Act 13/2003. Deca-sure: Reg. No. G4280 Act 36/1947, NS0 V19/24.4.2/1443 Act 13/2003. Levamintic Concentrate: Reg. No.G3569 Act 36/1947. NS0 V05/18.1.4/288 Act 13/2003. Levoxy 5.9%: Reg. No. G4148 Act 36/1947. Trio-sure: Reg. No. G4242 Act 36/1947, NS0 V18/24.4/1439 Act 13/2003

KAH/PM051/2023/1