


B-CO BOLIC

Injectable Tonic / Insputbare Tonikum

Reg. No./Nr. G4005 Act/Wet 36/1947
NSO V14/19.4/1245 Act 13/2003

Injectable tonic for Phosphorous, Selenium and Vitamin B deficiencies in ruminants, pigs and horses.

Insputbare tonikum vir Fosfor, Selenium en Vitamiene B tekorte in herkouers, varke en perde.



**INJECTABLE TONIC
INSPUTBARE TONIKUM**



B-CO BOLIC

Injectable solution

INDICATIONS

- Injectable tonic for Phosphorous, Selenium and Vitamin B deficiencies in ruminants, pigs and horses.
- To normalize and/or boost metabolism.
- To aid in the recovery after disease.

COMPOSITION per ml

Phosphorous	6 mg
Casein acid (digested)	50 mg
Selenium	0,01 mg
Vitamin B12	0,05 mg
Vitamin B1	1 mg
Vitamin B6	0,66 mg

STORAGE INSTRUCTIONS

Store between 5°C and 25°C

Protect from direct sunlight.

WARNINGS

- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.
- Do not inject intramuscularly.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Shake well before use and during administration.

Method**RUMINANTS:** Intravenous or subcutaneous injection.**HORSES and PIGS:** Subcutaneous injection only.**Dosage**

Cattle	
Adult	10 - 25 ml
Calves	5 - 15 ml
Horses	
Adult	15 - 20 ml
Ponies	5 - 15 ml

Sheep and goats	
Adult	3 - 5 ml
Lambs / kids	2 - 3 ml
Pigs	
Adult	5 - 10 ml
Piglets	3 - 5 ml

In healthy animals half of the above dose can be given. Do not depart from the recommended dosages without consulting an animal nutritionist or veterinarian.

B-CO BOLIC

Insipuitbare oplossing

AANDUIDINGS

- Insipuitbare tonikum vir Fosfor, Selenium en Vitamiene B tekorte in herkouers, varke en perde.
- Om metabolisme te normaliseer/verbeter.
- Om herstel na 'n siektetoestand te versnel.

SAMESTELLING per ml

Fosfor	6 mg
Kaseïen suur (verteer)	50 mg
Selenium	0,01 mg
Vitamiene B12	0,05 mg
Vitamiene B1	1 mg
Vitamiene B6	0,66 mg

BERGINGSAANWYSINGS

Berg tussen 5°C en 25°C

Beskerm teen direkte sonlig.

WAARSKUWINGS

- Hou buite bereik van kinders, oningesigte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.
- Moet nie binnespiers insipuit nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

Skud goed voor- en gedurende gebruik.

Metode**HERKOUERS:** Binneaars- of onderhuidse inspuiting.**PERDE en VARKE:** Slegs onderhuidse inspuiting.**Dosering**

Beeste	
Volwasse	10 - 25 ml
Kalwers	5 - 15 ml
Perde	
Volwasse	15 - 20 ml
Ponies	5 - 15 ml

Skape en bokke	
Volwasse	3 - 5 ml
Lammers / bokkies	2 - 3 ml
Varke	
Volwasse	5 - 10 ml
Varkies	3 - 5 ml

In gesonde diere kan helfte van die bogenoeme dosis gebruik word. Alvorens van aanbevole dosisse afgewyk word, moet 'n voedingskundige of veearts eers geraadpleeg word.