



For animal use only

③+⑧†

ENDO+LINT

Reg.No G3570 Act 36/1947
NSO V05/18.1.8/289 Act 13/2003

GIVE YOUR ANIMAL THE EDGE TO BEAT INTERNAL PARASITES

Double active combination ORAL dose remedy

Contains:

Levamisole HCl	37,5 mg / mℓ
Praziquantel	18,8 mg / mℓ



† Imidothiazole
† Isoquinoline

CAUTION

Store in a cool and dry place



Available in: 200 mℓ, 500mℓ, 1ℓ, 5ℓ & 10ℓ



Roundworm remedy



Tapeworm remedy

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF TREATMENT.
- No worm remedy should be dosed to sheep and goats that have not been immunised against pulpy kidney.
- When treated with levamisole, the odd animal may develop an allergic-type reaction of a transient nature which normally disappears within 24 hours. An antihistamine may be administered. Under no circumstances must cortisone preparations be used.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- Sheep and goats: 2 ml / 10 kg body mass.
- Cattle: 10 ml / 50 kg body mass.
- Ostriches: 1 ml / 1 kg body mass.

EFFICACY

Worm species	Sheep/goats		Cattle	
	Immature	Adult	Immature	Adult
Wireworm	*	*	*	*
Brown stomachworm		**		**
Long-neck bankruptworm	**	*		
Large-mouthed bowelworm	*	*		
Nodular worm	*	*	**	*
Lungworm	**	**		
Hookworm	*	*		*
Bankruptworm (<i>Trichostrongylus</i> spp.)	*	*		
Bankruptworm (<i>Cooperia</i> spp.)			*	*
Milk tapeworm	CLASS I			
	Ostriches			
Wireworm (<i>Libostrongylus douglassi</i>)			**	*
Hookworm (<i>Codiostomum struthionis</i>)			**	*
Tapeworm (<i>Houttuynia struthionis</i>)			**	*

* Controls ≥ 90 % effective

** Aids in control = 60 – 89 % effective

Class 1: 100 % effective in more than 80 % of treated animals.

This list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.