

Ivermax LA Platinum

Reg. No. G3832 Act 36/1947

[NSO] V13/18.1.2/1186 Act 13/2003



FOR ANIMAL USE ONLY

IVERMAX PLATINUM LA 
3,15 % Injectable solution

CAUTION

INDICATIONS

Antiparasitic remedy for cattle.

COMPOSITION

Ivermectin 3,15 % m/v

STORAGE INSTRUCTIONS

Store in a cool place, away from direct sunlight, in the original container tightly capped.

WARNINGS

- DO NOT SLAUGHTER CATTLE FOR HUMAN CONSUMPTION WITHIN 120 DAYS OF LAST TREATMENT
- DO NOT USE IN LACTATING CATTLE WHERE MILK OR MILK PRODUCTS ARE USED FOR HUMAN CONSUMPTION.
- DO NOT USE IN DAIRY CATTLE WITHIN 120 DAYS BEFORE CALVING.
- DO NOT ADMINISTER INTRAVENOUSLY OR INTRAMUSCULARLY.
- Keep out of reach of children, unformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Do not re-use empty containers.
- Refrain from smoking and eating while handling the product.
- Wash hands thoroughly after use.
- Appropriate vaccination precautions should be taken in areas where the risk of clostridial infections of cattle such as black quarter and malignant oedema are present.
- Studies indicate that when ivermectin comes in contact with the soil, it readily and tightly binds to the soil and become inactive over time. Drug containers and any residual contents should be disposed of by burying or incinerating as free ivermectin may adversely affect fish and certain water-borne organisms.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Dosage:

- 1 ml per 50 kg body mass.
- At the recommended dosage IVERMAX PLATINUM LA can be used in breeding animals as no detrimental reproductive effects have been reported

Dosage table	
Body mass (kg)	Dosage volume (ml)
Up to 50	1,0
51 – 100	2,0
101 – 150	3,0
151 – 200	4,0
201 - 250	5,0
251 – 300	6,0
301 – 350	7,0
351 – 400	8,0
401 – 450	9,0
451 - 500	10,0

Administration:

• SUBCUTANEOUS INJECTION ONLY

Note: Temporary discomfort has been observed in animals following subcutaneous injection. The pain reaction is sometimes intense, but usually transient. A low incidence of soft tissue swelling at the injection site has been observed. These reactions have disappeared without treatment.

- Inject cattle under the loose skin in front of or behind the shoulder.
- Use of a 2,0 cm 16 gauge needle is recommended.
- Use sterile equipment and follow aseptic procedures.
- Do not dilute or mix with any other substances.

EFFICACY: EXTERNAL PARASITES

- Kills sucking lice, biting lice and mange mites.
- Kills cattle screw-worms present at time of injection and protects against screw-worm strike for 2 weeks after administration.
- Controls sand tansans for up to 3 days after treatment.
- Controls blue ticks (*Rhipicephalus decoloratus* and *Rhipicephalus microplus*) on cattle for 89 days.

EFFICACY:

Cattle roundworms	Immature	Adult	Controls against re-infestation for
Wireworm (<i>Haemonchus placei</i>)	*	*	42 days
Brown Stomachworm (<i>Ostertagia ostertagi</i>)	*	*	49 days
Cattle Bankruptworm (<i>Cooperia</i> spp.)	*	*	42 days
Hookworm (<i>Bunostomum phlebotomum</i>)	*	*	
Nodular worm (<i>Oesophagostomum radiatum</i>)	*	*	49 days
Lungworm (<i>Dictyocaulus viviparus</i>)	*	*	63 days
Eyeworm (<i>Thelazia</i> spp.)		*	
<i>Parafilaria bovicola</i> (false bruising)		**	

* Control ≥ 90 % effective

** Aid in Control = 60 – 89 % effective

Instructions for use with automatic dosing equipment:

- Disinfect all needles and syringes before using by boiling in clean water for 15 to 20 minutes.
- Boiled needles should be stored in an antiseptic solution before use and changed frequently when injecting animals.
- Remove the draw-off assembly from the sterile pack. Handle carefully to avoid contamination.
- Connect the plastic tube firmly to dosing syringe. Use stepped adaptor provided if needed.
- Remove cap from bottle and disinfect rubber stopper with methylated spirits or other suitable chemical disinfectant. Hold bottle upright and fully insert draw-off needle into centre of rubber stopper.
- Hang bottle comfortably in inverted position from neck, shoulder or belt. Use attachment tapes provided.
- Gently prime injector. Equipment is now ready for use.
- After use, remove draw-off assembly from pack and flush out entire apparatus with water before storing. If connecting tube is re-used, it should be boiled for 15 to 20 minutes before use along with the injecting syringe and needles.
- Store partly used bottle in carton to protect from light.
- Do not re-use empty bottles.

† Macrocyclic lactone



Registration Holder: Kyrón Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.

KAH/INS/108/2022/1