

BE SURE. TRIO-SURE

Trio-Sure



PROTECTION AGAINST

- ANTHRAX
- BOTULISM
- BLACK QUARTER



250ML - 50 DOSES
50ML - 10 DOSES



250ML - 83 DOSES
50ML - 16 DOSES

Reg. No. G4242 Act 36/1947 **NSO** V18/24.4.2/1439 Act 13/2003

Available in:
50ML & 250ML



Registration Holder: Kyron Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal, Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.

FOR ANIMAL USE ONLY

VACCINE FOR CATTLE AND SHEEP

STORAGE INSTRUCTIONS

Store between 2 °C to 8 °C in a refrigerator. Do not freeze. Protect from light.

DOSAGE FORM

Injectable vaccine

COMPOSITION

Bacillus anthracis unencapsulated avirulent Sterne 34F₂ strain active spores, *Clostridium botulinum* type C₁ + C₂ and type D toxoids and *Clostridium chauvoei* anaerobes.

INDICATIONS

For the active immunisation of cattle and sheep against anthrax, botulism and blackquarter.

MODE OF ACTION

The active component stimulates the immune response against 3 diseases. Immunity is established within 3-4 weeks after inoculation, and lasts for at least one year.

WARNINGS

- Do not slaughter animals intended for human consumption within 21 days of last treatment.
- Keep out of reach of children, uninformed persons and animals.
- Usually no marked reaction follows vaccination although a transient swelling may appear at the site of inoculation, and some animals may show a moderate rise of temperature for one or two days.
- As for all vaccines, absolute immunity cannot be guaranteed in all animals. Therefore, it is advisable to consult a veterinarian regarding additional control measures.
- As with all vaccines, occasional hypersensitivity may occur. In such cases, appropriate treatment such as adrenaline and/or histamines should be administered without delay.
- Although this vaccine has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Concurrent administration of antibiotics to vaccinated animals is contraindicated as the antibiotic will interfere with the vaccine.
- Antibiotics should not be given for seven days before and after vaccination.

DIRECTIONS FOR USE

Use only as directed. Shake well before use. The recommended route of administration is by subcutaneous injection.

DOSAGE

Cattle: 5 ml

Sheep: 3 ml

DOSAGE RECOMMENDATIONS

- **CATTLE:** Calves should first be vaccinated at 4-6 months of age. A booster inoculation with a vaccine containing *Clostridium botulinum* type C₁ + C₂ and type D toxoids and *Clostridium chauvoei* anaerobes should be administered 4-6 weeks post initial vaccination. A single annual booster with Trio-Sure is recommended thereafter. For adult animals that have never been immunised before, vaccinate first with Trio-Sure and boost 4-6 weeks later with a vaccine containing *Clostridium botulinum* type C₁ + C₂ and type D toxoids and *Clostridium chauvoei* anaerobes. A single annual booster with Trio-Sure is recommended thereafter.
- **SHEEP:** Sheep should first be vaccinated at 3 - 4 months of age. A booster inoculation with a vaccine containing *Clostridium botulinum* type C₁ + C₂ and type D toxoids and *Clostridium chauvoei* anaerobes should be administered 4-6 weeks post initial vaccination. A single annual booster with Trio-Sure is recommended thereafter. For adult animals that have never been immunised before, vaccinate first with Trio-Sure. A booster inoculation with a vaccine containing *Clostridium botulinum* type C₁ + C₂ and type D toxoids and *Clostridium chauvoei* anaerobes should be administered 4-6 weeks post initial vaccination. A single annual booster with Trio-Sure is recommended thereafter, especially before any operation likely to produce wounds. In case of fleece it is necessary to ensure the needle has penetrated the skin before injecting the product.

GENERAL RECOMMENDATIONS

- Shake well before use.
- Inoculate by subcutaneous injection.
- Dispose of partly used containers.
- Use sterile needles and properly sterilised vaccinator. In order to maintain proper hygiene, it is advisable to change to clean sterile needles as frequently as possible.

PRESENTATION

50 ml & 250 ml.