



Reg. No. G4197 Act 36/1947 NSO V19/18.2/1442 Act 13/2003

# TICK + ROUNDWORM CONTROL

DOUBLE ACTIVE COMBINATION  
FIPRONIL 0,9 % + ABAMECTIN 0,5 %



- ✓ Ready-to-use Pour-on
- ✓ Residual action
- ✓ Control against Blue ticks
- ✓ Control against Multi-host ticks
- ✓ Registered against pyrethroid resistant blue ticks
- ✓ Broad-spectrum roundworm control



Registration Holder: Kyrón Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Roolhuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.

KAH/PM011/2022/1





Reg. No. G4197 Act 36/1947  
 [NSD] V19/18.2/1442 Act 13/2003

## FOR ANIMAL USE ONLY

## CAUTION

## COMPOSITION

Fipronil 0,9 % m/v  
 † Abamectin 0,5 % m/v

## INDICATIONS

A ready-to-use pour-on antiparasitic for cattle.

## STORAGE

Store at or below 25 °C, protected from sunlight.

## WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 105 DAYS OF LAST APPLICATION.
- DO NOT APPLY TO LACTATING ANIMALS IF THE MILK IS INTENDED FOR HUMAN CONSUMPTION.
- Do not apply to wet animals or if rain is expected shortly after application.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

## PRECAUTIONS

- Wear gloves during application.
- In case of accidental contact with the skin, wash with plenty of water.
- Do not eat, smoke or drink during application of the product.
- Do not re-use containers for any other purpose.
- Dispose of used packaging by burying or incineration.
- Avoid contamination of water sources.

## DIRECTIONS FOR USE:

USE ONLY AS DIRECTED

### Application

- Ready-to-use pour-on.
- Apply in two lines, on both sides of the spinal column, from the poll of the head to the tail root.

### Dosage

- 1 ml per 10 kg.

EFFICACY: EXTERNAL PARASITES	Controls for
Blue tick ( <i>Boophilus</i> spp.) including pyrethroid resistant strains	* 21 days ** 28 days
Red-legged tick ( <i>Rhipicephalus evertsi</i> )	* 21 days
Bont tick ( <i>Amblyomma</i> spp.)	* 14 days
Bont-legged tick ( <i>Hyalomma</i> spp.)	* 7 days ** 14 days
Brown ear tick ( <i>R.appendiculatus</i> )	* 7 days

\* Control ≥ 90 % effective

\*\* Aids in control = 60 – 89 % effective

EFFICACY: INTERNAL PARASITES	Immature	Adult
Brown stomach worm ( <i>Ostertagia</i> spp.)	*	*
Wireworm ( <i>Haemonchus placei</i> )	*	*
Hookworm ( <i>Bunostomum phlebotomum</i> )	*	*
Stomach bankruptworm ( <i>Trichostrongylus axei</i> )	*	*
Nodular worm ( <i>Oesophagostomum</i> spp.)	*	*
Cattle bankruptworm ( <i>Cooperia</i> spp.)	*	*

\* Control ≥ 90 % effective

KAH/PM011/2022/1