

ATTILA^{①†}

Reg. No. G4197 Act 36/1947

NS0 V19/18.2/1442 Act 13/2003

FOR EXTERNAL ANIMAL USE ONLY

CAUTION

COMPOSITION

Fipronil 0,9 % m/v
† Abamectin 0,5 % m/v

INDICATIONS

A ready-to-use pour-on antiparasitic for cattle.

STORAGE

Store at or below 25 °C, protect from sunlight.

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 105 DAYS OF LAST APPLICATION.
- DO NOT APPLY TO LACTATING ANIMALS WHERE THE MILK IS INTENDED FOR HUMAN CONSUMPTION.
- Do not apply to wet animals or if rain is expected shortly after application.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Wear gloves during application.
- In case of accidental contact with the skin, wash with plenty of water.
- Do not eat, smoke or drink during application of the product.
- Do not re-use containers for any other purpose.
- Dispose of used packaging by burying or incineration.
- Avoid contamination of water sources.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Application

- Ready-to-use pour-on.
- Apply in two lines, on both sides of the spinal column, from the poll of the head to the tail root.

Dosage

- 1 ml per 10 kg.

EFFICACY: EXTERNAL PARASITES	Controls for
Blue tick (<i>Boophilus spp.</i>) including pyrethroid resistant strains	* 21 days ** 28 days
Red-legged tick (<i>Rhipicephalus evertsi</i>)	* 21 days
Bont tick (<i>Amblyomma spp.</i>)	* 14 days
Bont-legged tick (<i>Hyalomma spp.</i>)	* 7 days ** 14 days
Brown ear tick (<i>R. appendiculatus</i>)	* 7 days

* Control ≥ 90% effective

** Aids in control = 60 – 89 % effective

EFFICACY: INTERNAL PARASITES	Immature	Adult
Brown stomach worm (<i>Ostertagia spp.</i>)	*	*
Wireworm (<i>Haemonchus placei</i>)	*	*
Hookworm (<i>Bunostomum phlebotomum</i>)	*	*
Stomach bankruptworm (<i>Trichostrongylus axei</i>)	*	*
Nodular worm (<i>Oesophagostomum spp.</i>)	*	*
Cattle bankruptworm (<i>Cooperia spp.</i>)	*	*

* Control ≥ 90% effective

† Macrocyclic lactone



Registration holder: Kyron Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.