

ENDOJECT + FLUKE^{®†}

Reg. No. G3648 Act 36/1947

FOR ANIMAL USE ONLY

INDICATIONS

Liver Fluke and Roundworm remedy for cattle with residual action against re-infestation of Wireworm, Hookworm and Nodular worm.

COMPOSITION

Closantel 7,5 % m/v.

STORAGE INSTRUCTIONS

Store in a cool place.

WARNINGS

- NOT FOR USE IN MILK COWS IF MILK IS INTENDED FOR HUMAN CONSUMPTION.
- ALLOW 30 DAYS BETWEEN LAST INJECTION AND SLAUGHTER FOR HUMAN CONSUMPTION.
- Handle with care. Poisonous by swallowing and inhalation.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Prevent contamination of food, food utensils, etc.
- Wash hands thoroughly after use.
- Use sterile injection apparatus. As for all injections a sharp sterile needle should be used for each animal.
- Do not inject animals whilst their coats are wet (e.g. after dipping) due to the risk of infection at the injection site.
- A transient swelling may occur at the injection site in some individual animals.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Administration

Can be used in all classes of cattle, irrespective of age, diet, condition, pregnancy, sex, breed or normal management practices.

By subcutaneous injection at the side of the neck or by intramuscular injection in any large muscle group using a 15 – 16 G needle 2,5 cm in length.

Do not exceed 20 ml per injection site.

Dosage:

- **Standard dose:** 1 ml / 30 kg body mass, e.g.:

| | | | | | | |
|------------------------|-----|-----|-----|-----|-----|-----|
| Body mass (kg) | 30 | 60 | 90 | 120 | 150 | 180 |
| Dose (ml) | 1,0 | 2,0 | 3,0 | 4,0 | 5,0 | 6,0 |
| No. of doses in 500 ml | 500 | 250 | 166 | 125 | 100 | 83 |

| | | | | | | |
|---------------------|-------|-----|-----|-----|-----|-----|
| Body mass (kg) | 30 | 60 | 90 | 120 | 150 | 180 |
| Dose (ml) | 1,0 | 2,0 | 3,0 | 4,0 | 5,0 | 6,0 |
| No. of doses in 1 l | 1 000 | 500 | 333 | 250 | 200 | 166 |

| | | | | | | |
|---------------------|-------|-------|-------|-------|-------|-----|
| Body mass (kg) | 30 | 60 | 90 | 120 | 150 | 180 |
| Dose (ml) | 1,0 | 2,0 | 3,0 | 4,0 | 5,0 | 6,0 |
| No. of doses in 5 l | 5 000 | 2 500 | 1 666 | 1 250 | 1 000 | 833 |

| | | | | | | |
|----------------------|--------|-------|-------|-------|-------|-------|
| Body mass (kg) | 30 | 60 | 90 | 120 | 150 | 180 |
| Dose (ml) | 1,0 | 2,0 | 3,0 | 4,0 | 5,0 | 6,0 |
| No. of doses in 10 l | 10 000 | 5 000 | 3 333 | 2 500 | 2 000 | 1 666 |

- **Double dose:** When a higher efficacy against internal parasites is indicated, a dose of 1 ml / 15 kg body mass can be administered, e.g.:

| | | | | | | |
|------------------------|-----|-----|-----|-----|------|------|
| Body mass (kg) | 30 | 60 | 90 | 120 | 150 | 180 |
| Dose (ml) | 2,0 | 4,0 | 6,0 | 8,0 | 10,0 | 12,0 |
| No. of doses in 500 ml | 250 | 125 | 83 | 62 | 50 | 41 |

| | | | | | | |
|---------------------|-----|-----|-----|-----|------|------|
| Body mass (kg) | 30 | 60 | 90 | 120 | 150 | 180 |
| Dose (ml) | 2,0 | 4,0 | 6,0 | 8,0 | 10,0 | 12,0 |
| No. of doses in 1 l | 500 | 250 | 166 | 125 | 100 | 83 |

| | | | | | | |
|---------------------|-------|-------|-----|-----|------|------|
| Body mass (kg) | 30 | 60 | 90 | 120 | 150 | 180 |
| Dose (ml) | 2,0 | 4,0 | 6,0 | 8,0 | 10,0 | 12,0 |
| No. of doses in 5 l | 2 500 | 1 250 | 833 | 625 | 500 | 416 |

| | | | | | | |
|----------------------|-------|-------|-------|-------|-------|------|
| Body mass (kg) | 30 | 60 | 90 | 120 | 150 | 180 |
| Dose (ml) | 2,0 | 4,0 | 6,0 | 8,0 | 10,0 | 12,0 |
| No. of doses in 10 l | 5 000 | 2 500 | 1 666 | 1 250 | 1 000 | 833 |

Efficacy

- **Standard dose:** 1 ml / 30 kg body mass:

| Liver Flukes | | | |
|---|---------|---------|-------|
| Liver fluke (<i>F. hepatica</i>) | 4 weeks | 6 weeks | Adult |
| | ** | ** | * |
| Giant liver fluke (<i>F. gigantica</i>) | 6 weeks | 8 weeks | Adult |
| | ** | ** | * |

| Roundworms | Immature | Adult |
|---|----------|-------|
| Wireworm (<i>Haemonchus</i> spp.) | * | * |
| Hookworm (<i>Gaigeria</i> spp.) | - | * |
| Nodular worm (<i>Oesophagostomum</i> spp.) | * | * |

| Residual efficacy | Interval between treatment and re-infestation |
|------------------------------------|---|
| Wireworm (<i>Haemonchus</i> spp.) | 3 weeks |

* Control ≥ 90 % effective.

** Aids in Control = 60 – 89 % effective.

Efficacy

- **Double dose:** 1 ml / 15 kg body mass:

| Liver Flukes | | | |
|---|---------|---------|-------|
| Liver fluke (<i>F. hepatica</i>) | 4 weeks | 6 weeks | Adult |
| | ** | ** | * |
| Giant liver fluke (<i>F. gigantica</i>) | 6 weeks | 8 weeks | Adult |
| | ** | * | * |

| Roundworms | Immature | Adult |
|---|----------|-------|
| Wireworm (<i>Haemonchus</i> spp.) | * | * |
| Hookworm (<i>Gaigeria</i> spp.) | * | * |
| Nodular worm (<i>Oesophagostomum</i> spp.) | * | * |

| Days where "control" is achieved | | |
|----------------------------------|----------|-------------|
| Wireworm | Hookworm | Nodularworm |
| 42 days | 21 days | 21 days |

* Control ≥ 90 % effective.

** Aids in Control = 60 – 89 % effective.

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.



Registration holder: Kyron Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooiwal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.