

# MOXIDOSE<sup>†</sup>

Reg. No. G4213 Act 36/1947

NSO V20/18.1.2/1445 Act 13/2003

## FOR ANIMAL USE ONLY

Oral antiparasiticide for sheep

**INDICATIONS:** For the treatment and control of roundworms and nasal worm in sheep.

## CAUTION

**STORAGE:** STORE BELOW 25 °C IN A COOL, DARK, DRY PLACE

**COMPOSITION:** Moxidectin 0.1 % m/v

## WARNINGS:

- DO NOT SLAUGHTER SHEEP FOR HUMAN CONSUMPTION WITHIN 14 DAYS OF LAST TREATMENT.
- Do not use in animals producing milk for human consumption.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

## PRECAUTIONS:

- Sheep should be vaccinated against pulpy kidney at least 3 weeks before treatment.
- Destroy empty containers and do not re-use for any other purpose.
- Ensure that aseptic injection technique and sterile equipment is used when administering the product.
- Do not eat, drink or smoke while handling the product.
- While using outside, store bottle in carton to protect from sunlight.
- Wash hands thoroughly after use.
- Do not dilute or mix with any other remedy

## USE ONLY ACCORDING TO THE FOLLOWING DIRECTIONS:

**Sheep:** Dose orally at 2 ml per 10 kg live mass

Dosage table	
Body mass (kg)	Dosage volume (ml)
Up to 10	2
11 – 20	4
21 – 30	6
31 – 40	8
41 – 50	10

## SAFETY:

Moxidose is safe at the recommended dosages. It may be used in sheep of all ages, including pregnant and breeding animals.

## EFFICACY:

Internal Parasites	Immature	Adult	Residual action <sup>+</sup>
<b>ROUND WORMS</b>			
Wire worm ( <i>Haemonchus contortus</i> )	*	*	28 days
Brown stomach worm ( <i>Teladorsagia circumcincta</i> )	*	*	28 days
Bankrupt worm ( <i>Trichostrongylus colubriformis</i> )	*	*	
Long-necked bankrupt worm ( <i>Nematodirus spathiger</i> )	*	**	
White bankrupt worm ( <i>Strongylus papillosus</i> )	*	X	
Hook worm ( <i>Gaigeria pachyscelis</i> )	*	*	28 days
Nodular worm ( <i>Oesophagostomum columbianum</i> )	*	*	28 days
Large-mouth bowel worm ( <i>Chabertia ovina</i> )	*	*	
Lung worm ( <i>Dictyocaulus filaria</i> )	*	*	

\* Controls = ≥ 90 % effective

\*\* Aids in control: 60% - 89% effective

+ Interval between treatment and re-infestation

X Ineffective

## Nasal Worm (*Oestrus ovis*):

MOXIDOSE controls nasal bot larvae (first, second and third instar larvae)

† Macrocylic lactone



Registration holder: Kyrón Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.