

Supatraz 25%

Reg. No. G3581 Act 36/1947
[NSO] V05/18.4.4/296 Act 13/2003

FOR EXTERNAL ANIMAL USE ONLY

Emulsifiable concentrate

OXPECKER COMPATIBLE

HARMFUL

STORAGE INSTRUCTIONS

Protect product from extremes of heat and cold and from dampness.

COMPOSITION

Amitraz 25,0% m/v

INDICATIONS

CATTLE: Controls ticks. Kills lice and mange mites.

SHEEP AND GOATS: Controls ticks. Controls sheep scab. Kills itch mites and goat mange mites.

WARNINGS

- WITHDRAWAL PERIOD: DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST DIPPING.
- DO NOT USE ON HORSES.
- In case of poisoning call a doctor and make this label available to him.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Tightly reseal the container if all the contents have not been used.
- Destroy empty containers.
- Wash hands thoroughly after use.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED

For external use on cattle, sheep and goats.

SUPATRAZ 25% should not be mixed with any other dip.

CATTLE

• Spraying

Dipwash should be freshly prepared for each spraying. Add required amount of SUPATRAZ 25% to \pm 10 ℓ water in a suitable container. When well mixed add to measured bulk water in spray sump and stir thoroughly. Use at a rate as per the table below:

SUPATRAZ 25%	WATER
100 m ℓ	100 ℓ
200 m ℓ	200 ℓ
500 m ℓ	500 ℓ
1 ℓ	1 000 ℓ
5 ℓ	5 000 ℓ
10 ℓ	10 000 ℓ
20 ℓ	20 000 ℓ

Boost

It is recommended that not more than 400 head be sprayed in the spray race at one time, especially where fouling is a problem. If, however, this figure is exceeded, boosting at the rate of 100 m ℓ per 200 head, commencing at 400 head, can be undertaken to maintain a satisfactory deposit of active ingredient, especially where tick-borne diseases are a problem.

• Plunge dip

Total replenishment: This requires that the diptank is topped up to the desired level on each dipping occasion. SUPATRAZ 25% is then added on every occasion at a rate of 1 ℓ to 5 000 ℓ water, taking the total amount of water in the tank into the calculation.

Lime stabilised:

- **Fresh fill:** Add SUPATRAZ 25% at the rate of 1 ℓ per 1 000 ℓ water. At the end of each dipping, add 6 kg calcium hydroxide (75%) per 1000 ℓ water.
- **Replenishment:** Add SUPATRAZ 25% at the rate of 1 ℓ per 1 000 ℓ water added. At the end of each dipping, calcium hydroxide (75%) must be added to the tank at a rate of 5 kg per 100 head of cattle dipped to ensure that the dipwash pH is in the region of 12.

Frequency:

- **Ticks and mange:** Spray/dip weekly
- **Lice:** Spray/dip twice with a 7 day interval

Should dipping intervals extend beyond 3 weeks, additional lime should be added at a rate of 1 kg lime (75% active) per 1 000 ℓ water in the diptank. The dipwash should be adequately protected for 2 months if this procedure is followed.

SHEEP AND GOATS

SUPATRAZ 25% should be administered by means of either:

1.2 Dippings using fresh, unstabilised dipwash. SUPATRAZ 25% should be added at a rate of 1 ℓ per 750 ℓ water, taking the total amount of water in the tank into consideration.

OR

2.A fresh, unstabilised dipwash for the first dipping, which is then stabilised with the addition of lime at a rate of 4 kg lime per 750 ℓ dipwash, added immediately after the initial dipping. This stabilised dipwash is then suitable for the second dipping, 7 days later.

Frequency:

Sheep and goats should be dipped twice with a 7 day interval to kill sheep scab, itch and goat mange mites. Ensure that the animals are immersed for approximately 1 minute with the heads being submerged at least 3 times in order to ensure that the inside of the ears are thoroughly wetted.

N.B.

Sheep scab is a notifiable disease and must be reported to the nearest State Veterinarian.

For the control of bacterial build-up and for the prevention of post-dipping lameness:

COPPER SULPHATE can be added at a rate of 500 g per 1 000 ℓ of water.

OR

ZINC SULPHATE can be added at a rate of 1 kg/1 000 ℓ of dipwash.

For the control of post-dipping lameness and lumpy wool:

ZINC SULPHATE can be added at a rate of 1 kg/400 ℓ of dipwash.

Registration Holder: Kyron Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.



Supatraz 25%

Reg. Nr. G3581 Wet 36 / 1947
[NSO] V05/18.4.4/296 Wet 13/2003

SLEGS VIR UITWENDIGE DIERGEGBRUIK

Emulsifiseerbare konsentraat

RENOSTERVOËL VERSOENBAAR

SKADELIK

BERGINGSAAWYSINGS

Beskerm teen uitermatige hitte en koue en teen klamheid.

SAMESTELLING

Amitras 25,0% m/v

INDIKASIES

BEESTE: Beheer bosluise. Dood luise en skurfte myte.

SKAPE EN BOKKE: Beheer bosluise. Beheer skaapbrandsiekte. Dood jeukmyte en bokskurfte myte.

WAARSKUWINGS

- ONTREKKINGSPERIODE: MOENIE DIERE VIR MENSLIKE VERBRUIK BINNE 7 DAE VANAF DIE LAASTE DIPPING SLAG NIE.
- MOENIE OP PERDE GEBRUIK NIE.
- In geval van vergiftiging raadpleeg 'n geneesheer en stel hierdie etiket aan hom beskikbaar.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

- Herseël die houer dig indien al die inhoud nie gebruik is nie.
- Vernietig leë houers.
- Was hande deeglik na gebruik.

GEBRUIKSAANWYSINGS – GEBRUIK SLEGS SOOS AANGEDUI

Vir uitwendige gebruik op beeste, skape en bokke.

SUPATRAZ 25% behoort nie met enige ander dip gemeng te word nie

BEESTE

• Bespuiting

Dipmengsel behoort vars opgemaak te word voor elke bespuiting.

Voeg die vereiste hoeveelheid SUPATRAZ 25% by ± 10 ℓ water in 'n geskikte houer. Wanneer dit deeglik gemeng is voeg dit by afgemete volume water in die dipmengselhouer en roer deeglik. Gebruik teen 'n koers soos in die tabel hieronder:

SUPATRAZ 25%	WATER
100 m ℓ	100 ℓ
200 m ℓ	200 ℓ
500 m ℓ	500 ℓ
1 ℓ	1 000 ℓ
5 ℓ	5 000 ℓ
10 ℓ	10 000 ℓ
20 ℓ	20 000 ℓ

Versterking

Dit word aanbeveel dat nie meer as 400 beeste op een tyd in die spuitdip bespuit word nie, veral waar besoedeling 'n probleem is. As hierdie getal oorskry word kan versterking teen 'n koers van 100 m ℓ per 200 diere, met aanvang van 400 diere, gedoen word om 'n aanvaarbare neerslag van aktiewe bestanddeel te verseker, veral waar bosluise oordraagbare siektes 'n probleem is.

• Dompeldip

Totale aanvulling: Dit verg dat die dipbak met elke dipping tot by die gewenste vlak aangevul word. SUPATRAZ 25% word dan met elke dipping bygevoeg teen 'n koers van 1 ℓ SUPATRAZ 25% by 5 000 ℓ water, met die totale hoeveelheid water in die dipbak in berekening gebring.

Stabilisering met kalk:

- **Varsvulling:** Voeg SUPATRAZ 25% by teen 'n koers van 1 ℓ per 1 000 ℓ water. Aan die einde van elke dipping, moet kalsium hidroksied (75%) by die dipbak gevoeg word teen 'n koers van 6 kg per 1000 ℓ water.
- **Aanvulling:** Voeg SUPATRAZ 25% by teen 'n koers van 1 ℓ per 1 000 ℓ water wat bygevoeg is. Aan die einde van elke dipping, moet kalsiumhidroksied (75%) by die dipbak gevoeg word teen 'n koers van 5 kg per 100 kop beeste gedip by die dipbak gevoeg word, om te verseker dat die dipmengsel se pH in die omgewing van 12 is.

Diptussenpose:

- **Bosluise en myte:** Bespuit/dip weekliks
- **Luise:** Bespuit/dip twee maal met 'n tussenpose van 7 dae

As die dip-tussenpose langer as 3 weke is, moet addisionele kalk bygevoeg word teen 'n koers van 1 kg kalk (75% aktief) per 1 000 ℓ water in die dipmengsel. As hierdie prosedure gevolg word behoort dit die dipmengsel vir 2 maande voldoende te beskerm.

SKAPE EN BOKKE

SUPATRAZ 25% behoort soos volg aangewend te word:

1.2 Dippings met vars, ongestabiliseerde dipmengsel. SUPATRAZ 25% moet bygevoeg word teen 'n koers van 1 ℓ per 750 ℓ water, met die totale hoeveelheid water in die dipbak in berekening gebring.

OF

2. 'n Vars, ongestabiliseerde dipmengsel vir die eerste dipping, wat dan gestabiliseer word met die byvoeging van kalk, teen 'n koers van 4 kg kalk per 750 ℓ dipmengsel, wat onmiddellik na die eerste dipping bygevoeg word. Hierdie gestabiliseerde dipmengsel is dan geskik vir die tweede dipping 7 dae later.

Diptussenpose:

Skaapbrandsiekte-, jeuk- en bokskurfte myte te dood. Verseker dat die diere vir ongeveer 1 minuut gedompel is, en dat hulle koppe ten minste 3 keer onderdompel word, om deeglike benatting van die binnekant van die ore te verseker.

L.W.

Skaapbrandsiekte is 'n aanmeldbare siekte en moet by die naaste Staatsveearts aangemeld word.

Vir die beheer van bakteriese opbouing en die voorkoming van nadipmankheid:

KOPERSULFAAT kan teen 'n koers van 500 g per 1 000 ℓ water bygevoeg word.

OF

SINKSULFAAT kan teen 'n koers van 1 kg/1 000 ℓ dipmengsel bygevoeg word.

Vir die beheer van nadipmankheid of klontwol:

SINKSULFAAT kan teen 'n koers van 1 kg/400 ℓ dipmengsel bygevoeg word.

Registrasiehouer: Kyron Animal Health (Edms) Bpk., Reg. Nr. 2004/021847/07, Eenheid 45C, 45 Parkview Straat, Highway Business Park, Rooihuiskraal Uitbr. 31, Centurion, 0157, Suid Afrika. Tel: 0861 247 463.