

Supatraz 125

Reg. No. G3011 Act 36/1947

[NSO] V14/18.3.4/1239 Act 13/2003

WARNINGS

DO NOT USE ON HORSES.

- Do not slaughter animals for human consumption within seven days of last dipping. (Milk from treated cows can be used for human consumption immediately after application.)
- Handle with care.
- Keep out of reach of children, uninformed persons and animals.
- In case of poisoning call a doctor and make this label available to him.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

Protect product from extremes of heat and cold and from dampness. Tightly reseal the container if all the contents have not been used. Destroy empty containers.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Only for spraying of cattle (hand spraying or spray races).

SUPATRAZ 125 should not be mixed with any other dip.

This dipwash must be freshly prepared for each spraying.

Mixing Instructions:

Premix the required amount of SUPATRAZ 125 with water in a suitable container. When well mixed, add the measured bulk water in the spray sump and stir thoroughly. Use at the rate of 2 parts SUPATRAZ 125 to 1 000 parts water as tabulated.

SUPATRAZ 125	WATER
200 ml	100 litre
2 litre	1 000 litre

Boost:

It is recommended that not more than 500 head be sprayed in the spray race at one time, especially where fouling is a problem. If however, this figure is exceeded, boosting at the rate of 200 ml per 100 head commencing at 500 head can be undertaken to maintain a satisfactory deposit of active ingredient, especially where tick-borne diseases are a problem.

Frequency of use: Intensive tick control:

When used as directed at weekly intervals, it will provide intensive tick control. All animals should be treated and interruption of the programme should be avoided as they will then, if not vaccinated, be susceptible to various tick-borne diseases.

Extended treatment intervals:

Under circumstances of low tick challenge, it may be used, at the user's discretion, at intervals greater than seven days. In these circumstances, users are advised to make use of relevant vaccines to obtain optimal protection of their cattle against prevalent tick-borne diseases.

Mange – spray weekly.

Lice – spray twice with a seven day interval.