

FOR ANIMAL USE ONLY / SLEGS VIR DIEREGERBUIK

TRINEX PLUS 19,5%^② + ^③



Broad spectrum anthelmintic drench for use in cattle, sheep & goats.

For the treatment and control of:

- ✓ Mature and immature gastrointestinal nematodes
- ✓ Lungworms
- ✓ Liver fluke

STORE BELOW 30 °C AND PROTECT FROM LIGHT.

CAUTION

Contains:

Levamisole 7,5 % m/v
Triclabendazole 12,0 % m/v

③ Imidothiazole ② Benzimidazole

Breë-spektrum wurmmiddel vir gebruik in beeste, skape en bokke.

Vir die beheer en behandeling van:

- ✓ Volwasse en onvolwasse maagderm rondewurms
- ✓ Longwurms
- ✓ Lewerslak

BERG BENEDE 30 °C EN BESKERM TEEN LIG.

VERSIGTIG

Bevat:

Levamisool 7,5 % m/v
Triklabendasool 12,0 % m/v



Registration Holder: Kyrón Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C,
45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.

Reg. No./Nr. G3814 Act/Wet 36/1947.
NSO V14/18.1.8/1244 Act/Wet 13/2003

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 35 DAYS OF LAST TREATMENT.
- DO NOT TREAT LACTATING COWS, SHEEP OR GOATS PRODUCING MILK FOR HUMAN CONSUMPTION.
- Milk cows, sheep and goats may be treated during the dry period.
- No worm remedy, not even TRINEX PLUS 19,5 %, should be dosed to sheep and goats that have not been immunised against pulpy kidney.
- Overdosing may cause transient side effects e.g. muscle tremors, salivation.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Clean drenching equipment before and after use.
- Do not contaminate dams, waterways or ditches with the product or used container.
- When administering, do not eat, drink or smoke.
- In case of accidental contamination rinse eyes and wash hands and skin immediately.
- In case of poisoning consult a doctor and make this label available.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED
SHAKE WELL BEFORE USE.
Do not dilute or mix with any other product.
Ready-to-use suspension.

Single oral dosage:

- **Cattle, sheep and goats**
1 ml TRINEX PLUS 19,5 % per 10 kg body mass.
1 ml is equivalent to 7,5 mg levamisole and 12 mg triclabendazole per kg body mass.

DOSAGE GUIDE

CATTLE		SHEEP & GOATS	
Body mass kg	Dosage ml	Body mass kg	Dosage ml
50	5	10	1
100	10	20	2
150	15	30	3
200	20	40	4
250	25	50	5
300	30	60	6
Each additional 50 kg	5	Each additional 10 kg	1

EFFICACY

CATTLE	Immature	Adult
Wireworm	*	*
Brown stomach worm	**	**
Bankrupt worm	*	*
Nodular worm	*	*
Hook worm		*
Liver fluke (<i>F. hepatica</i>)	All stages *	
Giant liver fluke (<i>F. gigantica</i>)	All stages *	
SHEEP & GOATS	Immature	Adult
Wireworm	*	*
Brown stomach worm	**	**
Bankrupt worm	*	*
Longnecked bankruptworm	*	*
Large-mouthed bankruptworm	*	*
Nodular worm	*	*
Hook worm		*
Liver fluke (<i>F. hepatica</i>)	All stages *	
Giant liver fluke (<i>F. gigantica</i>)	All stages *	

* Controls: ≥ 90 % effective

** Aids in control: 60 – 89 % effective

Treatment of animals suffering from worm infestations:

When animals are treated for worm infestations, all other animals with the same history should be treated as well. Possible subsequent treatment should be given according to veterinary advice. TRINEX PLUS 19,5 % is suitable for use with most drenching guns.

WAARSKUWINGS

- MOENIE DIERE VIR MENSLIKE VERBRUIK BINNE 35 DAE NA LAASTE BEHANDELING SLAG NIE.
- MOENIE LAKTERENDE KOEIE, SKAPE OF BOKKE WAT MELK VIR MENSLIKE VERBRUIK PRODUSEER, BEHANDEL NIE.
- Melkkoeie-, skape en bokke mag gedurende die droë periode behandel word.
- Geen wurmmiddel, nie eers TRINEX PLUS 19,5 %, moet aan skape en bokke gegee word wat nie teen bloednier geïmmuniseer is nie.
- Oordosering mag verbygaande nuwe effekte veroorsaak bv. spiertrillings, kwyling.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

DOSIS GIDS

BEESTE		SKAPE & BOKKE	
Liggaamsmassa kg	Dosis ml	Liggaamsmassa kg	Dosis ml
50	5	10	1
100	10	20	2
150	15	30	3
200	20	40	4
250	25	50	5
300	30	60	6
Elke addisionele 50 kg	5	Elke addisionele 10 kg	1

DOELTREFFENDHEID

BEESTE	Onvolwasse	Volwasse
Haarwurm	*	*
Bruinmaagwurm	**	**
Bankrotwurm	*	*
Knoppieswurm	*	*
Haakwurm		*
Lewerslak (<i>F. hepatica</i>)	Alle stadia *	
Reuse lewerslak (<i>F. gigantica</i>)	Alle stadia *	
SKAPE & BOKKE	Onvolwasse	Volwasse
Haarwurm	*	*
Bruinmaagwurm	**	**
Bankrotwurm	*	*
Langnekbankrotwurm	*	*
Grootbek bankrotwurm	*	*
Knoppieswurm	*	*
Haakwurm		*
Lewerslak (<i>F. hepatica</i>)	Alle stadia *	
Reuse lewerslak (<i>F. gigantica</i>)	Alle stadia *	

* Beheer: ≥ 90 % doeltreffend

** Hulpmiddel in beheer: 60 – 89 % doeltreffend

Behandling van diere met wurm besmettings:

Wanneer diere vir wurm besmettings behandel word, moet alle diere met dieselfde geskiedenis ook behandel word. Moontlike opvolgbehandelings moet volgens advies van 'n veearts toegedien word. TRINEX PLUS 19,5 % is geskik vir gebruik met meeste doseerspuitte.