

MAXISULF LA

Reg. No./Nr. G3657 Act/Wet 36/1947

NSO V14/17.1.7/1243 Act/Wet 13/2003



INJECTABLE SOLUTION / INSPUITBARE OPLOSSING

For the treatment of foot rot, navel-ill, joint-ill and pneumonia in stock, coccidiosis and bacterial scours in calves and lambs and strangles in horses.

Vir die behandeling van vrotpootjie, naelstringsiekte, septiese gewrigsontsteking en longontsteking by vee, koksidiose en bakteriese maagwerking by kalwers en lammers en nuwesiekte by perde.

CONTAINS / BEVAT

Sodium sulphadimethoxine 40 % mlv

Natriumsulfadimetoksien 40 % mlv



FOR ANIMAL USE ONLY
SLEGS VIR DIEREGERBUIK



Registration Holder: Kyrón Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.

STORAGE INSTRUCTIONS

Store in a cool dark place.

CONTAINS

Sodium sulphadimethoxine 40 % m/v

INDICATIONS

For the treatment of foot rot, navel-ill, joint-ill and pneumonia in stock, coccidiosis and bacterial scours in calves and lambs and strangles in horses.

- **Cattle, sheep and goats:**
Foot rot, pneumonia, navel-ill and joint-ill
- **Horses:**
Pneumonia, strangles, joint-ill and navel-ill
- **Pigs:**
Pneumonia and joint-ill.
- **Calves and lambs:**
Coccidiosis and bacterial scours

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST TREATMENT.
- MILK FROM TREATED COWS MUST NOT BE USED FOR HUMAN CONSUMPTION WITHIN 48 HOURS OF LAST TREATMENT.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE

USE ONLY AS DIRECTED

ADMINISTRATION

Administer by subcutaneous or intravenous routes. Massage the injection site to maximise absorption and to minimise swelling. When using the intravenous route, give the injection very slowly. Discontinue for a while at the first sign of discomfort and continue only when all seems normal.

N.B. Treated animals must have free access to ample drinking water during treatment and for at least two days after the last treatment.

DOSAGE

An initial dosage double that of the maintenance dosage is recommended. DO NOT TREAT FOR MORE THAN 4 DAYS.

AFTER ADMINISTERING THE INITIAL DOSE, ADMINISTER THE MAINTENANCE DOSE ONCE A DAY FOR 3 DAYS.

Species	Initial Dose	Maintenance Dose
Cattle, horses, sheep, goats and pigs	1 - 2 ml / 10 kg body mass	0,5 - 1 ml / 10 kg body mass

N.B. Consult your veterinarian Immediately if any uncertainty arises as to diagnosis or dosage or if little or no improvement is effected by the treatment.

BERGINGSAAWYSINGS

Berg in 'n koel, donker plek.

SAMESTELLING

Bevat Natriumsulfadimetoksien 40 % m/v

INDIKASIES

Vir die behandeling van vrotpootjie, naelstringsiekte, septiese gewrigsontsteking en longontsteking by vee, koksidiöse en bakteriese maagwerking by kalwers en lammers en nuwesiekte by perde.

- **Beeste, skape en bokke:**
Vrotpootjie, longontsteking, naelstringsiekte en septiese gewrigsontsteking.
- **Perde:**
Longontsteking, nuwesiekte, septiese gewrigsontsteking en naelstringsiekte.
- **Varke:**
Longontsteking en septiese gewrigsontsteking.
- **Kalwers en lammers:**
Koksidiöse en bakteriese maagwerking.

WAARSKUWINGS

- MOENIE DIERE BINNE 7 DAE NA DIE LAASTE BEHANDELING VIR MENSLIKE VERBRUIK SLAG NIE.
- MELK VAN BEHANDELDE KOEIE MOET NIE VIR MENSLIKE VERBRUIK BINNE 48 UUR NA LAASTE BEHANDELING GEBRUIK WORD NIE.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEBRUIKSAANWYSINGS

GEBRUIK SLEGS SOOS AANGEDUI

TOEDIENING

Dien toe deur onderhuidse- of binnearse inspuiting. Masseer die inspuitplek om opname te bevorder en swelling te verminder. Die binnearse inspuiting moet baie stadig toegedien word. Onderbreek die inspuiting by die eerste teken van ongemak en hervat inspuiting wanneer die dier normaal voorkom.

L.W. Behandelde diere moet vrye toegang tot volop drinkwater hê tydens behandeling en minstens twee dae na die laaste toediening.

DOSIS

'n Aanvangsdosis dubbel die hoeveelheid van die onderhoudsdosis word aanbeveel. MOET NIE VIR LANGER AS 4 DAE BEHANDEL NIE.

NADAT DIE AANVANKLIKE DOSIS TOEGEDIEN IS, DIEN DIE ONDERHOUDSDOSIS EEN MAAL 'N DAG VIR 3 DAE TOE.

Diersoort	Aanvangsdosis	Onderhoudsdosis
Beeste, perde, skape, bokke en varke	1 - 2 ml / 10 kg liggaamsmassa	0,5 - 1 ml / 10 kg liggaamsmassa

L.W. Indien enige onsekerheid ten opsigte van die dosis of diagnose ontstaan, of as geen beterskap intree na die behandeling nie, moet u onmiddelik u veearts raadpleeg.