

# MAXISULF LA

Reg. No. G3657 Act 36/1947

NSO VI4/17.1.7/1243 Act 13/2003

## STORAGE INSTRUCTIONS

Store in a cool dark place.

## COMPOSITION

Sodium sulphadimethoxine 40 % *m/v*

## INDICATIONS

For the treatment of foot rot, navel-ill, joint-ill and pneumonia in stock, coccidiosis and bacterial scours in calves and lambs and strangles in horses.

## WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST TREATMENT.
- MILK FROM TREATED COWS MUST NOT BE USED FOR HUMAN CONSUMPTION WITHIN 48 HOURS OF LAST TREATMENT.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder,

## DIRECTIONS FOR USE - USE ONLY AS DIRECTED

### Administration

Administer by subcutaneous or intravenous routes. Massage the injection site to maximise absorption and to minimise swelling. When using the intravenous route, give the injection very slow. Discontinue for a while at the first sign of discomfort and continue only when all seems normal. **N.B.** Treated animals must have free access to ample drinking water during treatment and for at least two days after the last treatment.

## INDICATIONS

- **Cattle, sheep and goats:**  
Foot rot, pneumonia, navel-ill and joint-ill
- **Horses:**  
Pneumonia, strangles, joint-ill and navel-ill.
- **Pigs:**  
Pneumonia and joint-ill
- **Calves and lambs:**  
Coccidiosis and bacterial scours.

## DOSAGE

An initial dosage double that of the maintenance dosage is recommended. DO NOT TREAT FOR MORE THAN 4 DAYS. **AFTER ADMINISTERING THE INITIAL DOSE, ADMINISTER THE MAINTENANCE DOSE ONCE A DAY FOR 3 DAYS.**

Species	Initial dose	Maintenance dose
Cattle, horses, sheep, goats and pigs	1 - 2 ml / 10 kg body mass	0,5 - 1 ml / 10 kg body mass

**N.B.** Consult your veterinarian immediately if any uncertainty arises as to diagnosis or dosage or if little or no improvement is effected by the treatment.