

MAXITET 10%

Reg. No. G3857 Act 36/1947

[NSO] V13/17.1.2/1199 Act 13/2003

STORAGE INSTRUCTIONS

Store in a cool, dark place.

As with all oxytetracycline solutions darkening may occur during use without affecting the efficacy.

INDICATIONS

For the treatment of heartwater, tick-borne gallsickness (anaplasmosis) pneumonia, footrot, joint-ill and navel-ill in cattle, sheep and goats.

COMPOSITION

A sterile, stable solution containing 100 mg Oxytetracycline per ml.

WARNINGS

- TREATED ANIMALS MUST NOT BE SLAUGHTERED FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST TREATMENT.
- MILK FROM TREATED ANIMALS MUST NOT BE USED FOR HUMAN CONSUMPTION WITHIN 48 HOURS OF LAST TREATMENT.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Livestock: Subcutaneous, intramuscular or intravenous injection.

Dosage: 1 ml / 10 kg body mass.

Notes:

Massage the area after injection to enhance absorption and minimise swelling.

• Intramuscular injection

Inject in the muscle of the mid-neck area with a maximum volume of 20 ml at any one site.

• Intravenous injection

Using a short needle inject very slowly and pause if any discomfort occurs.

Daily doses should be divided i.e. ½ of the daily dose should be administered at 12 hourly intervals.

• Repetition of dosages

Anaplasmosis and heartwater

Repeat treatment daily as required.

All other conditions

Treat daily for 3 – 4 days.

N.B. For rapid recovery a prompt and accurate diagnosis is essential. It is, therefore, always advisable to consult your veterinarian.



Registration Holder: Kyron Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.

MAXITET 10%

Reg. Nr. G3857 Wet 36/1947

NSO V13/17.1.2/1199 Wet 13/2003

BERGINGSANWYSINGS

Bêre in 'n koel, donker plek.

Soos met alle oksitetrasiklien oplossings mag verdonkering gedurende gebruik voorkom sonder om die doeltreffendheid te affekteer.

AANWYSINGS

Vir die behandeling van hartwater, bosluisoorgedraagde galsiekte (anaplasmose), longontsteking, vrotpootjie, naelstringontsteking en septiese gewrigsontsteking by beeste, skape en bokke.

SAMESTELLING

'n Steriele, stabiele oplossing wat 100 mg Oksitetrasiklien per ml bevat.

WAARSKUWINGS

- BEHANDELDE DIERE MOENIE BINNE 7 DAE NA LAASTE BEHANDELING VIR MENSLIKE VERBRUIK GESLAG WORD NIE.
- MELK VAN BEHANDELDE DIERE MOENIE VIR MENSLIKE VERBRUIK BINNE 48 URE NA BEHANDELING GEBRUIK WORD NIE.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

Lewendehawe: Onderhuidse, binnespiers of binnearse inspuiting.

Dosis: 1 ml / 10 kg liggaamsmassa.

Notas:

Masseer die inspuitplek om opname te bevorder en swelling te verminder.

• Binnespiers inspuiting

Spruit in die dik spier van die middelnekarea met 'n maksimum dosis van 20 ml per area

• Binnearse inspuiting

Gebruik 'n kort naald en spruit stadig met 'n tydelike onderbreking as ongemak ontstaan. Daaglikse dosisse moet verdeel word d.w.s. ½ van die daaglikse dosis moet met 12 uur tussenposes toegedien word.

• Herhaling van dosisse

Anaplasmose en hartwater

Herhaal behandeling daaglik soos benodig.

Alle ander toestande

Behandel daaglik vir 3 – 4 dae.

N.B. Vir vinnige herstel is 'n spoedige en akkurate diagnose noodsaaklik. Dit is dus altyd raadsaam om u veearts te raadpleeg.

