

FOR ANIMAL USE ONLY

STORAGE

Store at or below 25 °C. Protect from sunlight.

COMPOSITION

Each ml contains Sulphamethoxazole 200 mg, Trimethoprim 40 mg.

INDICATIONS

Maxi-Trim is indicated for use in cattle, sheep and pigs to treat sulphonamide sensitive infections. It is used to treat bacterial infections of the respiratory tract, gastrointestinal tract and urinary tract as well as wound infections and footrot.

WARNINGS

Withdrawal:

Meat: 5 days, Milk: 3 days.

- May cause swelling at the site of injection.
- Do not use in animals with liver disease.Do not use in animals with known sulpha drug sensitivity.
- Keep out of reach of children, animals and uninformed persons.
- Although this remedy has been extensively tested under a wide variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DOSAGE AND DIRECTIONS FOR USE

- Use only as directed.
- By intramuscular injection 1 ml per 15 kg body mass daily for 5 days. In cases of severe infection the dose can be increased to 1,5 ml per 15 kg body mass.

IDENTIFICATION

Yellow, slightly viscous transparent solution.

PRESENTATION

Amber glass vial containing 100 ml sterile solution.

