

FOR ANIMAL USE ONLY

# CALBRO FORTE

**INJECTION**

Reg. No. G4349 Act 36/1947

NSO V22/19.2/1501 Act 13/2003

## INJECTABLE SOLUTION

### INDICATIONS

It is indicated for hypocalcaemia and hypomagnesaemia – in cattle, sheep, pigs, and horses - as well as for ketosis in cattle.

### STORAGE INSTRUCTIONS

STORE IN ITS ORIGINAL CONTAINER IN A COOL AND DRY PLACE BELOW 30 °C. PROTECT FROM LIGHT.

### COMPOSITION

Each 100 ml contains:

Calcium Gluconate Monohydrate	40,0 g
eq. to 3,57 g elemental Calcium	
Magnesium Glycerophosphate	1,0 g
Magnesium Chloride Hexahydrate	3,0 g
eq. to 0,16 g elemental Phosphorus	
eq. to 0,48 g elemental Magnesium	
Anhydrous Glucose	5,0 g
Anhydrous Caffeine	0,4 g

### WARNINGS

- WITHDRAWAL PERIOD: ZERO DAYS FOR MILK AND MEAT
- For subcutaneous, intraperitoneal or intravenous administration.
- The intravenous administration can produce phlebitis or coagulation in the injection site. If injected very fast, heart arrhythmia and/ or bradycardia may result.
- Keep out of reach of children, uniformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### PRECAUTIONS

- In serious cases, the slow intravenous route is recommended combined with subcutaneous route. This ensures a quick initial effect followed by a more sustained effect. It is recommended that the product be gently heated to body temperature before being administered intravenously.
- When the injections are administered exclusively by subcutaneous route, the dose must be distributed in three or four sites. Intravenous injections must be administered slowly.
- Do not re-use empty containers.
- Once used, destroy packaging and surplus in the proper manner, according to local regulations.

### DIRECTIONS FOR USE: USE ONLY AS DIRECTED

#### Route of administration

Subcutaneous, intraperitoneal or intravenous.

#### Dosage:

- Bovine and Equine: 60 - 100 ml
- Calf and foal: 15 - 30 ml
- Ovine: 10 - 20 ml
- Swine: 10 - 30 ml

### PRESENTATION

Blow moulded, PP (Polypropylene) veterinary bottle with butyl stopper and safety aluminium seal (flip off).



*Registration Holder: Kyrón Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa, Tel: 0861 247 463.*

**Botswana Listing No.: BV2200611/A**  
**Category of Distribution: VPS**

KAH/INS016/2022/2