

FOR ANIMAL USE ONLY / SLEGS VIR DIEREGERBUIK

Doxy-Max 50 %

Reg. No./Nr. G3918 Act/Wet 36/1947
[NSZ] V14/17.1.2/1254 Act/Wet 13/2003

Water Soluble Antibiotic powder

For the treatment of bacterial respiratory disease and bacterial gastrointestinal infections in fowls.

Contains: Doxycycline hyclate 500 mg/g

Water Oplosbare Antibiotika Poeier

Vir die behandeling van bakteriële- asemhaling siektes en maagderm infeksies in hoenders.

Bevat: Doksisisiklien hiklaat 500 mg/g



Registration holder / Registrasiehouer: Kyron Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa.
Tel: 0861 247 463.

Doxy-Max 50 %

Reg. No./Nr. G3918 Act/Wet 36/1947
[NSZ] V14/17.1.2/1254 Act/Wet 13/2003

Water soluble powder

For the treatment of bacterial respiratory disease and bacterial gastrointestinal infections in fowls.

Storage instructions: STORE BELOW 25 °C

Contains: Doxycycline hyclate 500 mg/g

WARNINGS

- DO NOT SLAUGHTER FOR HUMAN CONSUMPTION WITHIN 9 DAYS AFTER LAST TREATMENT.
- DO NOT ADMINISTER TO HENS LAYING EGGS FOR HUMAN CONSUMPTION.
- Medicated drinking water should be consumed within 24 hours.
- Hypersensitivity reactions, gastrointestinal disturbances, hepatotoxicity and haematologic effects have been reported but are rare.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Skin contact with the product should be avoided.
- Keep lid tightly closed after the seal has been broken.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- For oral administration in drinking water.

Dosage

40 mg Doxy-Max 50 % (20 mg doxycycline) per kg body mass per day for 3 - 4 consecutive days.

Dosage table

- **1-19 day old broilers:** 0,1 g Doxy-Max 50 % per 1 ℓ drinking water
- **20-24 day old broilers:** 0,15 g Doxy-Max 50 % per 1 ℓ drinking water
- **25-30 day old broilers:** 0,2 g Doxy-Max 50 % per 1 ℓ drinking water

For smaller numbers of broilers:

- **1-19 day old broilers:** 1 level measuring spoon of 4 g / 5 ml Doxy-Max 50 % per 40 liter drinking water
- **20-24 day old broilers:** 1 level measuring spoon of 4 g / 5 ml Doxy-Max 50 % per 25 liter drinking water
- **25-30 day old broilers:** 1 level measuring spoon of 4 g / 5 ml Doxy-Max 50 % per 20 liter drinking water

These dosages can be affected by factors like ambient temperature, salt content in food, health status, stocking density, quality and flow rate of water.

Water oplosbare poeier

Vir die behandeling van bakteriese asemhaling siektes en maagderm infeksies in hoenders.

Bergingsaanwysings: BERG ONDER 25 °C

Bevat: Doksisisiklien hikklaat 500 mg/g

WAARSKUWINGS

- MOET NIE VIR MENSLIKE VERBRUIK BINNE 9 DAE NA LAASTE BEHANDELING SLAG NIE.
- MOET NIE TOEDIEN AAN HENNE WAT EIERS LÊ VIR MENSLIKE VERBRUIK NIE.
- Behandelde drinkwater moet binne 24 ure ingeneem word.
- Hipersensitieweits reaksies, maagderm ongesteldheid, hepatotoksiese- en hematologiese effekte is al aangemeld, maar is ongewoon.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid toestande getoets is, kan dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

- Vel kontak met die produk moet vermy word.
- Hou dig geseël nadat houer oopgemaak is.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- Vir orale toediening in drinkwater

Dosering

40 mg Doxy-Max 50 % (20 mg doksisisiklien) per kg liggaamsmassa per dag vir 3 - 4 opeenvolgende dae.

Doseringstabel

- **1-19 dae oud braaikuikens:** 0,1 g Doxy-Max 50 % per 1 ℓ drinkwater
- **20-24 dae oud braaikuikens:** 0,15 g Doxy-Max 50 % per 1 ℓ drinkwater
- **25-30 dae oud braaikuikens:** 0,2 g Doxy-Max 50 % per 1 ℓ drinkwater

Vir kleiner hoeveelhede hoenders:

- **1-19 dae oud braaikuikens:** 1 gelykvul maatlepel van 4 g / 5 ml Doxy-Max 50 % per 40 liter drinkwater
- **20-24 dae oud braaikuikens:** 1 gelykvul maatlepel van 4 g / 5 ml Doxy-Max 50 % per 25 liter drinkwater
- **25-30 dae oud braaikuikens:** 1 gelykvul maatlepel van 4 g / 5 ml Doxy-Max 50 % per 20 liter drinkwater

Hierdie doserings kan deur faktore beïnvloed word soos omgewings temperatuur, sout inhoud van voer, gesondheid, bevolkings digtheid, kwaliteit en vloei van water.