

Ovi Dose 4^{①③④⑧†}

Reg. No. G4041 Act 36/1947

NSO V14/18.1.8/1247 Act 13/2003

FOR ANIMAL USE ONLY

INDICATIONS

Oral drench for sheep and goats.

COMPOSITION

Praziquantel	2,50 % m/v
Levamisole	4,00 % m/v
Closantel	3,75% m/v
Ivermectin	0,10 % m/v
Sodium selenite	0,11 % m/v
As Selenium base	0,97 mg/mℓ
Cobalt chloride	0,10 % m/v
As Cobalt base	0,70 mg/mℓ

STORAGE INSTRUCTIONS

Store below 25°C. Protect from light.

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 49 DAYS OF LAST TREATMENT.
- Vaccination against clostridial infections e.g. pulpy kidney (enterotoxaemia), at least 3 weeks prior to treatment with OVI DOSE 4 is recommended.
- When treated with levamisole, an animal may occasionally develop an allergic-type reaction of a transient nature which normally disappears within 24 hours. An antihistamine may be administered. Do not administer cortisone preparations. Consult a veterinarian.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Do not contaminate water sources with product or containers.
- Do not re-use empty containers and dispose of container and residual contents in accordance with waste disposal regulations.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED SHAKE WELL BEFORE USE

- Ensure that the product is not dosed into the lungs.
- Ensure dosing equipment is clean before use. Clean all equipment after use.

DOSAGE

Dose orally at 1 mℓ per 5 kg

Dosage table	
Body mass (kg)	Dosage volume (mℓ)
15	3.0
20	4.0
25	5.0

30	6.0
35	7.0
40	8.0
45	9.0
50	10.0
55	11.0
60	12.0

EFFICACY

Internal parasites

ROUNDWORMS	Immature	Adult
Wireworm (<i>Haemonchus contortus</i>)	*	*
Bankruptworm (<i>Trichostrongylus colubriformis</i>)	*	*
Long-necked bankruptworm (<i>Nematodirus spathiger</i>)	*	*
White bankruptworm (<i>Strongyloides papillosus</i>)	*	*
Large-mouth bowelworm (<i>Chabertia ovina</i>)	*	*
Brown stomachworm (<i>Teladorsagia circumcincta</i>)	*	*

Nasal worm (<i>Oestrus ovis</i>)	*	*
---------------------------------------	---	---

Residual Efficacy	Interval between treatment and reinfestation
Wireworm (<i>Haemonchus contortus</i>)	28 days

TAPEWORM	
Milk tapeworm (<i>Moniezia expansa</i>)	Class I
Serrated tapeworm (<i>Thysanezia giardi</i>)	Class I

FLUKES	Immature	Adult
Conical fluke (<i>Calicophoron</i> spp)	*	
Liver fluke (<i>F. hepatica</i>)	*	*
Giant liver fluke (<i>F. gigantica</i>)	*	*

* Controls ≥ 90 % effective

† ① Macroyclic lactone, ③ Imidothiazole, ④ Salicylanilide, ⑧ Isoquinoline



Registration holder: Kyrón Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.