


# MAXI-BCO SUPER

Reg. No. G4543 Act/Wet 36/1947

FOR ANIMAL USE ONLY / SLEGS VIR DIEREGEBRUIK

A HIGH POTENCY VITAMIN B  
COMPLEX INJECTION  
'N HOË STERKTE VITAMIEN B  
KOMPLEKS INSPUITING



Registration Holder / Registrasiehouer: Kyron Animal Health (Pty) Ltd., Reg. No. 2004/021847/07, Unit 45C, 45 Parkview Street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa. Tel: 0861 247 463.



## MAXI-BCO SUPER

Injectable solution

For the supportive therapy and treatment of stress and B Vitamin deficiencies in all animals.

### COMPOSITION:

#### Each ml contains:

Vitamin B <sub>1</sub>	50 mg
Vitamin B <sub>2</sub>	7,5 mg
Vitamin B <sub>6</sub>	10 mg
Vitamin B <sub>12</sub>	100 mcg
Nicotinamide	100 mg
Pantothenic acid	10 mg

### DOSAGE AND DIRECTIONS FOR USE:

**Route of administration:** Intramuscular or subcutaneous injection

Dogs & Cats:	0,5 - 2 ml
Pigs & Sheep:	5 - 10 ml
Cattle & Horses:	20 - 30 ml

### STORAGE:

Store in a cool dark place at or below 25 °C.  
Shake well before use.

### WARNING:

Keep out of reach of children and uninformed persons.

In healthy animals half of the above dose can be given. Do not depart from the recommended dosages without consulting an animal nutritionist or veterinarian.

## MAXI-BCO SUPER

Inspuitbare oplossing

Vir die ondersteunende terapie en behandeling van stres en Vitamien B tekorte by alle diere.

### SAMESTELLING:

#### Elke ml Bevat

Vitamiën B <sub>1</sub>	50 mg
Vitamiën B <sub>2</sub>	7,5 mg
Vitamiën B <sub>6</sub>	10 mg
Vitamiën B <sub>12</sub>	100 mcg
Nikotienamied	100 mg
Pantoteensuur	10 mg

### DOSERING EN GEBRUIKSAANWYSINGS:

**Roete van toediening:** Binnespiere- of onderhuidse inspuiting

Honde & Katte:	0,5 - 2 ml
Varke & Skape:	5 - 10 ml
Beeste & Perde:	20 - 30 ml

### BERGINGSANWYSINGS:

Bewaar in 'n koel, donker plek by of onder 25 °C. Skud goed voor gebruik.

### WAARSKUWING:

Hou buite bereik van kinders en oningeligte persone.

In gesonde diere kan helfte van die bogenoemde dosis gebruik word. Alvorens van aanbevole dosisse afgewyk word, moet 'n voedingskundige of veearts eers geraadpleeg word.

