

For Animal Use Only

## **Curaclox DC** Intramammary

Reg. No. G 2617 (Act 36/1947)

### **Storage:**

Store in a cool place.

### **Composition:**

Each 4,5 g syringe contains 500 mg cloxacillin (as the benzathine salt) and 250 mg ampicillin (as trihydrate) in a long-acting base.

### **Indications For Use:**

Curaclox DC is indicated for the prevention and treatment of mastitis during the dry period.

### **Warnings:**

Milk for human consumption may only be taken from 60 hours post calving (i.e. 28 days plus 60 hours after treatment).

Cattle may be slaughtered for human consumption only after 28 days from last treatment.

Not to be used in lactating cows. Do not use within 28 days before calving.

Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### **Directions for Use:**

Use only as directed.

After final milking of a lactation, clean and disinfect the teats.

Introduce the contents of one syringe into each quarter via the teat canal.

### **Registration Holder:**



iPharma (Pty) Ltd  
(Reg: 2010/000192/07)  
124 Elevation Avenue,  
Randjesfontein, Midrand, 1683  
[www.ipharma.co.za](http://www.ipharma.co.za)

### **Distributed by:**



Kyrón Animal Health (Pty) Ltd  
(Reg. No. 2004/021847/07)  
Unit 45C, 45 Parkview street, Highway  
Business Park, Rooihuiskraal Ext. 31,  
Centurion, 0157.  
Tel: 0861 247 463

Namibia: V00/16.1/265