

For Animal Use Only

Norcal P.M.D.

Reg. No. G2869 (Act 36/1947)

Indications:

For the treatment of hypocalcaemia complicated by deficiency of magnesium with accompanying hypoglycaemia in cattle, horses, sheep, pigs and goats.

Storage:

Store below 25 °C. Protect from light.

Composition:

Norcal P.M.D. is a clear pale yellow solution containing calcium borogluconate 20 % m/v, dextrose 22 % m/v, magnesium hypophosphite 5 % m/v and 0,1 % chlorocresol as preservative.

Warnings:

Withdrawal period for meat or milk is nil. Keep out of reach of children, uninformed persons and animals. Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Precautions:

The solution should be warmed to body temperature before administration. Intravenous injections should be given slowly, in order to avoid possible neuromuscular dysfunction and cardiac insufficiency.

Normal aseptic procedures should be observed. For subcutaneous or slow intravenous injection.

Directions for Use:

Use only as directed.

By subcutaneous or slow intravenous injection.

Cattle: 200 - 400 ml

Horses: 150 - 300 ml

Sheep, pigs and goats: 15 - 50 ml

Registration Holder:



iPharma (Pty) Ltd

Reg. No.: 2010/000192/07

124 Elevation Avenue, Randjesfontein,
Midrand, 1683

www.ipharma.co.za

Marketed and Distributed by:



KYRON
AGRI

Kyrón Animal health (Pty) Ltd

Reg. No. 2004/021847/07

Unit 45C, 45 Parkview Street, Highway
Business Park, Rooihuiskraal Ext. 31,
Centurion, 0157

Tel: 0861 247 463

Slegs vir dieregebruik

Norcal P.M.D.

Reg. Nr. G2869 (Wet 36/1947)

Indikasies:

Vir die behandeling van hipokalsemie wat gekompliseer word deur 'n magnesiumtekort met gepaardgaande hipoglisemie, in beeste, perde, skape, varke en bokke.

Berging:

Berg benede 25 °C. Beskerm teen lig.

Samestelling:

Norcal P.M.D. is 'n helder liggeel oplossing wat kalsium boroglukonaat 20 % m/v, dekstrose 22 % m/v, magnesiumhipofosfiet 5 % m/v, en 0,1 % kloorkresol as 'n preserveermiddel, bevat.

Waarskuwings:

Daar is geen onttrekkingsperiode vir vleis of melk nie. Hou buite die bereik van kinders, ongeligte persone en diere. Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

Voorsorgmaatreëls:

Die oplossing behoort na liggaamstemperatuur opgewarm te word voor toediening. Binnearse inspuitings behoort stadig gegee te word om moontlike neuromuskulere afwykings en kardiaale ontoereikendheid te voorkom.

Normale aseptiese voorsorgmaatreëls moet gehandhaaf word. Vir onderhuidse of stadige binnearse inspuiting.

Gebruiksaanwysings:

Gebruik slegs soos aangedui. Vir onderhuidse of stadig binnearse inspuiting.

Beeste: 200 - 400 ml

Perde: 150 - 300 ml

Skape, varke en bokke: 15 - 50 ml

Registrasiehouer:



iPharma (Edms) Bpk
(Reg. No. 2010/000192/07)

Elevationlaan 124,
Randjesfontein, Midrand, 1683
www.ipharma.co.za

Versprei en Bemerk deur:



Kyrón Animal Health (Edms) Bpk
Reg. Nr. 2004/021847/07
Unit 45C, 45 Parkview straat,
Highway Business Park, Rooihuiskraal
Ext. 31, Centurion, 0157.
Tel: 0861 247 463