

DRY COW HEALTH PROGRAM



Optional vaccinations*

E.coli, Rota- & Corona Virus, Salmonella, Lumpy Skin Disease (LSD), Rift Valley Fever (RVF), Three Day Stiff sickness (Bovine Ephemeral Fever)

Internal & External Parasite Control



OR

Liver fluke treatment - all stages



OR

Trace mineral and Vitamin A & E supplement. For optimal immunity, udder health, colostrum & fertility.*



Oil based injectable vitamin A, D₃ and E supplement



Vaccinate to control clostridial diseases + Botulism + Anthrax



28 - 55 Days
Average = 36 Days

15 - 25 Days
Average = 21 Days


DRY-OFF

STEAM-UP



BREEDING

CALVING



Calving to Conception
Average = 90 Days

Voluntary Waiting Period
Average = 60 Days

Supplement for the treatment of milk fever & ketosis



Supplement for the treatment of milk fever & ketosis



MILK PRODUCTION

COLOSTRUM PRODUCTION

Dry cow intramammary



Optional vaccinations*

Campylobacteriosis, Trichomoniasis, Brucellosis, Respiratory & Reproductive Viruses (BVDV/IBR/BRSV/PI3), Mannheimia, Pasteurella



SCAN HERE FOR PRODUCT CATALOGUE

* Optional vaccinations based on disease prevalence and risk – Consult your veterinarian for recommendations.

*This program guideline is used entirely at your own risk and the registration holder does not accept any liability due to any products used and / or actions implemented.

Registration Holder: Kyron Animal Health (Pty) Ltd., Reg. No. 2004/021847/07,


Unit 45C, 45 Parkview street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa, Tel: 0861 247 463.

References:

1. SPEARS, J.W. & WEISS, W.P. 2008. Role of antioxidants and trace elements in health and immunity of transition dairy cows, *The Veterinary Journal* 176: 70-76
2. VAN EMON, M., SANFORD, C. & MCCOSKI, S. 2020. Review: Impacts of Bovine Trace Mineral Supplementation on Maternal and Offspring Production and Health, *Animals*, 10: 2404; doi:10.3390/ani10122404

MEDICATION USE & DOSE

Reg. No. C3582 Act 36/1947
NSO V05/18.12/293 Act 13/2003




Antiparasitic remedy.

Cattle
Injection (subcutaneous) **1 ml per 50 kg**

Milk withdrawal: Do not use in lactating cattle where milk or milk products are used for human consumption. Do not use in dairy cattle within 28 days of calving.

Reg. No. C2617 Act 36/1947
NSO V0016/265 Act 13/2003




Antibiotic treatment & prevention of mastitis during the dry period.

Cattle
Intramammary **1 syringe per teat**

Milk withdrawal: Milk for human consumption may only be taken from 60 hours post calving (i.e. 28 days plus 60 hours after treatment). Do not use in lactating cows. Do not use within 28 days before calving.

Reg. No. C3571 Act 36/1947
NSO V05/18.12/290 Act 13/2003




Dip controlling ticks and flies. Kills lice and protects against screw-worm.

Cattle
Spray Race, Hand Spray

Spray race: 1,5 l per 1000 l water
Hand spraying: 15 ml per 10 l water

Milk withdrawal: Zero days

Reg. No. C4349 Act 36/1947
NSO V22/19.2/1501 Act 13/2003




Treatment of hypocalcaemia (milk fever), hypomagnesaemia and ketosis.

Cattle
Injection (subcutaneous or intravenous)

Adult: 60 - 100 ml
Calf: 15 - 30 ml

Milk withdrawal: Zero days

Reg. No. C3619 Act 36/1947
NSO V14/18.11/238 Act 13/2003




Liver fluke remedy against immature & adult liver fluke and giant liver fluke.

Cattle
Oral Dose **12 ml per 100 kg**

Milk withdrawal: 14 days

Reg. No. C2869 Act 36/1947




Treatment of hypocalcaemia (milk fever) complicated by deficiency of magnesium with accompanying hypoglycaemia.

Cattle
Subcutaneous or Slow Intravenous **200 - 400 ml**

Milk withdrawal: Zero days

Reg. No. C3814 Act 36/1947
NSO V14/18.11/244 Act 13/2003




Roundworm and liver fluke (all stages) remedy.

Cattle
Oral Dose **5 ml per 50 kg**

Milk withdrawal: Do not treat lactating cows producing milk for human consumption. Milk cows may be treated during the dry period.


Reg. No. C4120 Act 36/1947
NSO V17/19.2/1390 Act 13/2003



Injectable trace mineral (Se-Met, Cu, Mn, Zn) and vitamin A & E solution.

Cattle
Injection (subcutaneous) **1 ml per 50 kg**


Reg. No. C4280 Act 36/1947
NSO V19/24.4.2/1443 Act 13/2003



Vaccine for immunisation against 10 important *Clostridium* strains.

Cattle
Injection (subcutaneous) **5 ml per animal**

Reg. No. C3796 Act 36/1947
NSO V14/19.12/41 Act 13/2003




Injectable vitamin A, D₃ & E solution.

Cattle
Injection (subcutaneous or into muscle)

Weaner: 0,5 ml
Yearling: 1 ml
Adult: 2 ml


Reg. No. C4258 Act 36/1947
NSO V18/24.4/1436 Act 13/2003



Vaccine for immunisation against Anthrax and Botulism

Cattle
Injection (subcutaneous) **4 ml per animal**

Reg. No. C4005 Act 36/1947
NSO V14/19.4/1245 Act 13/2003



Injectable tonic to boost metabolism and aid in recovery after disease.

Cattle
Injection (subcutaneous).

Adult: 10 - 25 ml
Calf: 5 - 15 ml

TREATMENT FOR CALVES WITH DIARRHOEA

DAY 1:
Inject Maxi-Trim.
Dose Diomec Plus Powder.



DAY 2:
Inject Maxi-Trim.
Dose Diomec Plus Powder.
Inject B-co Bolic.



DAY 3 - 5 (as necessary):
Inject Maxi-Trim.
Dose Diomec Plus Powder.



In case of roundworm and / or tapeworm infection – dose Endo + Lint



Company Representative:

Contact Number:

Registration Holder: Kyrón Animal Health (Pty) Ltd., Reg. No. 2004/021847/07,
Unit 45C, 45 Parkview street, Highway Business Park, Rooihuiskraal Ext. 31, Centurion, 0157, South Africa, Tel: 0861 247 463.